



EVANGELISTA ORTHOPEDIC CLINIC

6740 East Camelback Road, Suite 101
Scottsdale, AZ 85251
480-656-0291

Gregory T. Evangelista, M.D.

Amanda Damiris, PA-C

Rotator Cuff Repair Post-Operative Rehabilitation Guideline

0-6 Weeks:

- Goal: Establish Passive Range of Motion (PROM)¹
- Exercise: PROM ¹ONLY (No pulleys, No wands)
- Sling:
 - Wear sling at all times
 - If you move frequently at night, you should wear your sling while you sleep.
- Note: If a subscapularis repair was performed, please limit Exercise range (ER) to 15 degrees for the first 6 weeks

6-12 Weeks:

- Goal: Begin strengthening
- Exercises: Initiate Active Assisted Range of Motion (AAROM)¹
Begin with pendulum, pulley, and wand exercises
- Sling: As needed. May recommend in large crowds to avoid re-injury

12+ Weeks:

- Goal: Increase Strength
- Exercises: Continue Active Range of Motion (AROM)* and begin progressive strengthening (low load, high repetition therapeutic exercise), initiate wall climbing

¹ All PROM, AAROM, and AROM is to be performed in the plane of the scapula (forward elevation) or with the arm adducted (internal and external rotation)

Note:

- For the first 6 weeks, external rotation is limited to 30 degrees and internal rotation is limited to the chest wall
- AROM/AAROM for internal and external rotation should always be performed with the arm adducted
- No abduction is permitted for the first 6 weeks. Passive abduction is permitted thereafter

If you have any questions about these guidelines, your procedure or the rehabilitation process please do not hesitate to call us at **480-656-0291**