

EVANGELISTA ORTHOPEDIC CLINIC

Gregory T. Evangelista, M.D.

Perry J. Evangelista, M.D. 3271 N. Civic Center Plaza, Suite 110 Scottsdale, AZ 85251

Phone: (480) 656-0291 Fax: (480) 656-0127

ROTATOR CUFF REPAIR POST-OPERATIVE REHABILITATION GUIDELINES

0-6 Weeks:

- Goal: Establish Passive Range of Motion (PROM)¹
- Exercise: PROM ¹ONLY (No pulleys, No wands)
- Sling:
 - o Wear sling at all times
 - o If you move frequently at night, you should wear your sling while you sleep.

Note: If a subscapularis repair was performed, please limit Exercise range (ER) to 15 degrees for the first 6 weeks

6-12 Weeks:

- Goal: Begin strengthening
- Exerc Initiate Active Assisted Range of Motion (AAROM)¹
 - Begin with pendulum, pulley, and wand exercises
- Sling: As needed. May recommend in large crowds to avoid re-injury

12+ Weeks:

- Goal: Increase Strength
- Exerc Continue Active Range of Motion (AROM)* and begin progressive strengthening (low load, high repetition therapeutic exercise), initiate wall climbing

¹ All PROM, AAROM, and AROM is to be performed in the plane of the scapula (forward elevation) or with the arm adducted (internal and external rotation)

Note:

- For the first 6 weeks, external rotation is limited to 30 degrees and internal rotation is limited to the chest wall
- AROM/AAROM for internal and external rotation should always be performed with the arm adducted
 - No abduction is permitted for the first 6 weeks. Passive abduction is permitted thereafter

If you have any questions about these guidelines, your procedure or the rehabilitation process please do not hesitate to call us at 480-656-0291.